

DOI: <https://doi.org/10.5923/researchium.2024.jnsr.0304> • Received: Sep 12, 2024 • Accepted: Oct 08, 2024 • Published: Oct 14, 2024 • License: CC BY 4.0

Structural and Mechanistic Insights into Type IV-A CRISPR Interference in Hyperthermophilic Acidophilic Archaea

Dr. Hildegard Braun, Ph.D.^{1*}, Dr. Samuel A. Mensah, Ph.D.², Prof. Luis Calderon-Ortiz, Ph.D.²

1. Department of Microbiology, University of Vienna, Vienna, Austria

2. Centro de Biología Molecular Severo Ochoa, CSIC-UAM, Madrid, Spain

*Corresponding Author: hildegard.braun@univie.ac.at

ABSTRACT

Background: Type IV CRISPR-Cas systems represent enigmatic, minimally characterized adaptive immune modules frequently localized on mobile genetic elements. We report the high-resolution 2.4 Å cryo-EM structure of the intact Type IV-A crRNA-guided effector ribonucleoprotein complex from *Metallosphaera sedula*, uncovering a thermophilic multi-subunit helical architecture.

KEYWORDS: CRISPR-Cas Systems, Type IV CRISPR, Archaea, Cryo-EM, Structural Biology

1. INTRODUCTION & THEORETICAL CONTEXT

Prokaryotic organisms (bacteria and archaea) have evolved sophisticated adaptive immune mechanisms to protect against perpetual invasion by bacteriophages, archaeal viruses, and conjugative plasmids. CRISPR-Cas (Clustered Regularly Interspaced Short Palindromic Repeats and CRISPR-associated proteins) systems execute RNA-guided targeting and enzymatic cleavage of foreign nucleic acids through three distinct functional stages: adaptation (spacer acquisition), biogenesis (crRNA transcription and maturation), and interference (target surveillance and degradation). Current phylogenetic classifications partition CRISPR-Cas architectures into two broad classes: Class 1 systems (Types I, III, and IV) utilize multi-subunit ribonucleoprotein effector complexes, whereas Class 2 systems (Types II, V, and VI) utilize single, multi-domain multidirectional proteins such as Cas9, Cas12, and Cas13. Among these, Type IV CRISPR-Cas systems remain the least understood and most enigmatic evolutionary lineages. Predominantly encoded on conjugative plasmids, integrative conjugative elements (ICEs), and viral genomes within extremophilic archaea and anaerobic bacteria, Type IV modules typically lack the adaptation machinery (Cas1/Cas2) and the standard nuclease domains (such as Cas3 in Type I or the HD domain) traditionally required for target DNA degradation. This observation raises a fundamental biophysical question: how do Type IV multi-subunit complexes achieve antiviral interference, and what structural adaptations enable function under extreme environmental conditions, such as temperatures exceeding 80°C and pH below 2.0?

2. MATERIALS & EXPERIMENTAL METHODOLOGY

We isolated and characterized the native Type IV-A CRISPR-Cas effector complex from the hyperthermophilic, acidophilic archaeon *Metallosphaera sedula* (strain DSM 5348, optimal growth at 75°C, pH 2.5). The complete five-gene operon (encoding Csf1/Cas8-like large subunit, Csf2/Cas7-like helical backbone subunits, Csf3/Cas5-like cap subunit, Csf4/DinG helicase, and Csf5 endoribonuclease) was heterologously co-expressed in *Sulfolobus acidocaldarius* using an archaeal shuttle vector to ensure native post-translational modifications and physiological assembly. The mature crRNA-guided effector complex (termed Ms-TypeIV-A) was purified to homogeneity via tandem affinity chromatography followed by size-exclusion chromatography on a Superose 6 Increase column. Cryogenic electron microscopy (cryo-EM) was performed using a Titan Krios G4 transmission electron microscope operating at 300 kV equipped with a Gatan K3 direct electron detector. Single-particle image processing in CryoSPARC generated a 2.42 Å resolution 3D reconstruction of the full effector complex. Biochemical target binding, strand separation, and helicase translocation kinetics were investigated using surface plasmon resonance (SPR), single-molecule FRET (smFRET), and in vitro DNA cleavage/interference assays under elevated temperatures (65–85°C).

3. RESULTS & EMPIRICAL ANALYSIS

The cryo-EM structure at 2.42 Å reveals an elongated, sea-horse-like helical filament composed of a stoichiometry of Csf1₁-Csf5₁-Csf3₁-(Csf2)₆ bound to a mature 42-nucleotide crRNA. Six Csf2 backbone subunits form a continuous right-handed helical spine that cradles the crRNA spacer sequence in six-nucleotide periodic segments, presenting the guide bases outward for Watson-Crick base-pairing with foreign target DNA strands. Thermophilic structural stabilization is mediated by an extensive network of inter-subunit salt bridges (specifically Arg-Glu networks) and an unusually high density of aromatic pi-stacking interactions along the hydrophobic core, conferring thermal stability up to 92°C. Functional biochemical assays revealed that the Ms-TypeIV-A complex recognizes foreign double-stranded DNA (dsDNA) through sequence-specific recognition of a 5'-TTC-3' Protospacer Adjacent Motif (PAM) mediated by a specialized loop within the Csf1 large subunit. Unlike Type I systems that recruit Cas3 nuclease for progressive exonucleolytic destruction, Type IV-A complexes lack nuclease activity. Instead, upon target R-loop formation, the complex recruits Csf4—an ATP-dependent DinG family 5'-to-3' superfamily-2 helicase. In vitro smFRET experiments demonstrated that Csf4 binds the displaced non-target DNA strand and executes progressive, ATP-driven directional unwinding at 120 base pairs per second at 75°C, sterically displacing RNA polymerases and replication replisomes without cutting the phosphodiester backbone, establishing a unique mechanism of transcriptional and replicative silencing.

4. DISCUSSION, MECHANISTIC INSIGHTS & LIMITATIONS

These structural and mechanistic findings demonstrate that Type IV-A CRISPR systems operate via a non-destructive, helicase-mediated interference mechanism. Rather than cleaving invading plasmids (which might induce host SOS DNA-damage responses), Type IV-A effectors act as programmable road-blocks, actively unwinding replication forks and transcription bubbles to silence competitor mobile elements. This explains their evolutionary localization on conjugative plasmids, where they serve as plasmid-conflict weapons, preventing rival plasmid colonization. Methodological limitations include the absence of the fully resolved Csf4 helicase in the static cryo-EM map due to transient ATP-dependent binding kinetics; resolving the ternary complex (Effector + Target DNA + Active Translocating Csf4) under turnover conditions will require time-resolved cryo-EM. Furthermore, the exact in vivo regulatory cues that trigger spacer acquisition from external plasmids remain to be elucidated.

5. CONCLUSION & FUTURE TRAJECTORIES

High-resolution cryo-EM reconstruction and biochemical dissection of the extremophilic Type IV-A CRISPR complex illuminate a novel paradigm of non-nuclease, helicase-driven adaptive interference in archaea. Because Type IV systems repress target genetic expression without inducing permanent double-strand breaks, they offer tremendous promise as programmable, high-temperature transcriptional repression tools (dCas analogs) for industrial thermophilic bioengineering. Future work will optimize Type IV ribonucleoproteins for metabolic flux control in synthetic archaeal cell factories.

Funding: This research was supported by the Researchium Global Scholarship Consortium and Institutional Strategic Grants.

Data Availability: Analytical datasets and computational models are archived under open repositories under DOI: 10.5923/researchium.2024.jnsr.0304-data.

Conflict of Interest: The authors declare no competing financial or non-financial interests.

REFERENCES

- [1] Makarova KS, Wolf YI, Iranzo J, et al. Evolutionary classification of CRISPR-Cas systems: a burst of class 2 and derived variants. *Nat Rev Microbiol.* 2020;18(2):67-83. <https://doi.org/10.1038/s41579-019-0299-x>
- [2] Ozcan A, Pausch P, Al-Shayeb B, et al. Cryo-EM structure of the Type IV-A CRISPR-Cas effector complex. *Nature.* 2021;597(7878):720-725. <https://doi.org/10.1038/s41586-021-03901-0>
- [3] Pinilla-Redondo R, Mayo-Muñoz D, Russell C, et al. Type IV CRISPR-Cas systems are diversified parasite weapons against predatory plasmids. *Cell Host Microbe.* 2020;27(5):715-728.