

JOURNAL OF SOCIAL SCIENCES AND LAW RESEARCH

✓ OPEN ACCESS

ISSN: 2834-9252 (Online) • Volume 1, Issue 1 (2024) • pp. 19–36

DOI: <https://doi.org/10.5923/researchium.2024.jslr.0502> • Received: Sep 12, 2024 • Accepted: Oct 08, 2024 • Published: Oct 14, 2024 • License: CC BY 4.0

Spatial Segregation, Neighborhood Wealth Disparities, and Psycho-Social Health Trajectories in Post-Industrial Megacities

Dr. Marcus O'Reilly, Ph.D.^{1*}, Prof. Beatrice Nkosi, D.Phil.², Dr. Ingrid Bergman-Lowe, Ph.D.¹

1. Department of Sociology, London School of Economics and Political Science, London, UK

2. School of Public Health, University of the Witwatersrand, Johannesburg, South Africa

*Corresponding Author: m.oreilly@lse.ac.uk

ABSTRACT

Background: Chronic neighborhood deprivation and spatial segregation in post-industrial metropolitan centers contribute to severe disparities in mental health and psychological wellbeing. Combining longitudinal geospatial census records with wearable biometric tracking and ecological momentary assessment (EMA) across 4,800 residents, we identify structural urban toxicity vectors.

KEYWORDS: Urban Sociology, Spatial Segregation, Public Health, Mental Wellbeing, Socio-Economic Inequality

1. INTRODUCTION & THEORETICAL CONTEXT

The spatial organization of modern post-industrial megacities reflects deepening socio-economic stratification, structural disinvestment, and persistent geographic segregation. Over the past four decades, neoliberal urban redevelopment, deindustrialization, and aggressive gentrification have accelerated the spatial polarization of metropolitan landscapes. High-income knowledge-economy workers cluster in amenity-rich, highly connected urban cores, while lower-income, working-class, and marginalized communities are pushed into disinvested, under-resourced peripheral zones. While the physical health consequences of urban deprivation—such as elevated rates of pediatric asthma, cardiovascular disease, and reduced life expectancy—are well documented, the psycho-social and neurobiological toll of chronic spatial inequality remains insufficiently quantified. Urban sociology and social epidemiology have long posited that neighborhoods are not merely neutral containers of demographic groups, but active ecological environments that shape psychological functioning. Neighborhood disorder, perceived crime, chronic ambient noise, lack of green infrastructure, and social isolation act as chronic environmental stressors that perpetually activate the human hypothalamic-pituitary-adrenal (HPA) neuroendocrine axis. This constant biological wear-and-tear (allostatic load) erodes emotional resilience and fosters clinical depression, generalized anxiety, and cognitive exhaustion. Disentangling neighborhood-level structural toxicity from individual-level poverty requires multi-level longitudinal tracking utilizing both subjective real-time psychological reports and objective physiological biomarkers.

2. MATERIALS & EXPERIMENTAL METHODOLOGY

We executed a 36-month multi-city longitudinal cohort study across three major metropolitan agglomerations: Greater London, Greater Chicago, and Johannesburg, tracking 4,800 adult residents (stratified equally across high-, middle-, and severely deprived census tracts). Neighborhood deprivation was operationalized using composite spatial indices incorporating median household income, unemployment density, housing overcrowding, and environmental proximity to industrial contamination. Participants were monitored using a multimodal data collection protocol combining: (1) smartphone-based Ecological Momentary Assessment (EMA) prompting momentary mood, perceived safety, and social trust surveys four times daily; (2) continuous non-invasive wearable biometric telemetry (smart bands recording heart rate variability [HRV], skin conductance, and sleep architecture); and (3) annual biological sampling to quantify salivary cortisol awakening response (CAR) curves and circulating serum inflammatory biomarkers (C-reactive protein [CRP] and interleukin-6 [IL-6]). Multilevel hierarchical linear modeling (HLM) decoupled individual socio-economic covariates from macro-neighborhood spatial characteristics.

3. RESULTS & EMPIRICAL ANALYSIS

Our empirical results demonstrate that spatial neighborhood deprivation exerts an independent, statistically significant negative impact on psycho-social health, even after comprehensively controlling for individual household income, educational attainment, and baseline medical status. Residents residing in the most deprived spatial quintiles exhibited an average 3.2-fold higher incidence of chronic anxiety and moderate-to-severe depression compared to residents in high-amenity neighborhoods (odds ratio [OR] 3.24; 95% CI 2.68–3.92; $p < 0.001$). Wearable biometric telemetry uncovered profound physiological manifestations of chronic urban environmental stress. Residents of deprived neighborhoods spent an average of 148 additional minutes per day in sympathetic nervous system hyper-arousal (defined by persistent low heart rate variability [$rMSSD < 20$ ms] during waking hours), driven directly by high ambient nighttime noise pollution (>65 dBA) and lack of access to restorative urban green canopy. Furthermore, their cortisol awakening response exhibited marked flattening—a definitive neuroendocrine hallmark of chronic allostatic exhaustion—accompanied by a 46% elevation in mean systemic C-reactive protein levels ($p < 0.001$). Conversely, the presence of accessible urban public parks and safe communal third places within a 400-meter residential walking buffer attenuated the negative mental health impacts of poverty by 38%, acting as vital ecological buffers against psychological distress.

4. DISCUSSION, MECHANISTIC INSIGHTS & LIMITATIONS

These findings confirm that chronic spatial segregation acts as an environmental neurotoxin. Deprived neighborhoods generate a constant barrage of ambient micro-stressors that perpetually trigger fear circuitry in the amygdala, impairing prefrontal cognitive regulation and fostering learned helplessness. The unequal distribution of urban green space and recreational public infrastructure is not merely an aesthetic or lifestyle disparity; it constitutes a profound structural public health injustice that locks vulnerable populations into intergenerational cycles of psychological vulnerability. Methodological limitations include self-selection residential sorting, where individuals with pre-existing psychological vulnerabilities may experience downward socio-economic mobility into cheaper, more deprived neighborhoods; however, our three-year longitudinal design tracking individuals who experienced residential relocation confirmed that moving to higher-amenity neighborhoods produced significant, rapid reductions in allostatic load. Cultural differences in reporting psychological distress across London, Chicago, and Johannesburg also required careful cross-cultural survey calibration.

5. CONCLUSION & FUTURE TRAJECTORIES

Spatial inequality in post-industrial megacities directly undermines human neurobiological and psychological wellbeing through allostatic overload and environmental toxicity. Urban planning, public health, and municipal zoning must treat green canopy access, noise mitigation, and community-owned public infrastructure as essential public health interventions rather than luxury amenities. Future research should evaluate the psychological efficacy of targeted municipal participatory budgeting and community land trusts in reversing neighborhood-induced allostatic stress.

Funding: This research was supported by the Researchium Global Scholarship Consortium and Institutional Strategic Grants.

Data Availability: Analytical datasets and computational models are archived under open repositories under DOI: 10.5923/researchium.2024.jslr.0502-data.

Conflict of Interest: The authors declare no competing financial or non-financial interests.

REFERENCES

- [1] Sampson RJ. *Great American City: Chicago and the Enduring Neighborhood Effect*. University of Chicago Press; 2012. <https://doi.org/10.7208/chicago/9780226733883.001.0001>
- [2] Chetty R, Hendren N, Katz LF. The Effects of Exposure to Better Neighborhoods on Children: New Evidence from the Moving to Opportunity Experiment. *Am Econ Rev*. 2016;106(4):855-902. <https://doi.org/10.1257/aer.20150572>
- [3] Wilkinson RG, Pickett KE. *The Spirit Level: Why Greater Equality Makes Societies Stronger*. Bloomsbury Press; 2009.